

DO-TH 2005/01

January 2005

Radiatively Generated Isospin Violations in the Nucleon

M. Glück, P. Jimenez-Delgado, E. Reya

*Universität Dortmund, Institut für Physik,
D-44221 Dortmund, Germany*

Abstract

Isospin violating valence and sea distributions are evaluated due to QED leading $\mathcal{O}(\alpha)$ corrections to the standard QCD evolution equations. Unique perturbative predictions are obtained within the radiative parton model, and confirm earlier results. Nonperturbative contributions have been estimated and depend on a single free parameter chosen to be the current quark mass. The relevance of our predictions for extracting $\sin^2 \theta_W$ from DIS $\nu(\bar{\nu})N$ data ('NuTeV anomaly') is discussed as well.

The NuTeV collaboration recently reported [1] a measurement of the Weinberg angle $s_W^2 \equiv \sin^2 \theta_W$ which is approximately three standard deviations above those presented in [2] for the world average of other electroweak measurements. Possible sources for this discrepancy (see, for example, [3, 4, 5, 6, 7]) include, among other things, an isospin violating contribution δR_I^- to the relation [8] $R_{PW}^- = \frac{1}{2} - s_W^2$,

$$R^- \equiv \frac{\sigma_{NC}^{\nu N} - \sigma_{NC}^{\bar{\nu} N}}{\sigma_{CC}^{\nu N} - \sigma_{CC}^{\bar{\nu} N}} = \frac{1}{2} - s_W^2 + \delta R_I^- . \quad (1)$$

This contribution is given, for $N = \frac{1}{2}(p+n)$, by [3, 5, 7]

$$\delta R_I^- = \left(\frac{1}{2} - \frac{7}{6} s_W^2 \right) \frac{\delta U_v - \delta D_v}{U_v + D_v} \quad (2)$$

where the second moments $\delta Q_v(Q^2)$ and $Q_v(Q^2)$ of the valence distributions $(\delta)q_v(x, Q^2)$ are $Q_v(Q^2) = \int_0^1 x q_v(x, Q^2) dx$, $q_v = q - \bar{q}$, and $\delta Q_v(Q^2) = \int_0^1 x \delta q_v(x, Q^2) dx$ with

$$\begin{aligned} \delta u_v(x, Q^2) &= u_v^p(x, Q^2) - d_v^n(x, Q^2) \\ \delta d_v(x, Q^2) &= d_v^p(x, Q^2) - u_v^n(x, Q^2) . \end{aligned} \quad (3)$$

NLO QCD corrections do not significantly [4, 6, 7] contribute to (1). Nonperturbative calculations [9, 10, 11] of δq_v in (3) were found [5] to possibly reduce the discrepancy between the neutrino and electroweak measurements of s_W^2 by up to 40%. A somewhat different calculation [12], based on QED contributions to $\delta q_v(x, Q^2)$, resulted in a comparable reduction of this discrepancy.

We shall approach the issue of the isospin violations within the framework of the radiative parton model [13] and obtain predictions for δq_v which depend on a single free parameter required by the nonperturbative contribution and chosen to be, as in [12], the current quark mass. Although our method differs from those in [5], the results turn out to be similar. On the other hand, our starting position is only a little different to [12], the results, as anticipated in that work, turn out to be quite similar.

Following [12] we shall evaluate the modifications of the parton distributions due to QED radiative effects. In contrast to [12, 14] we shall calculate these effects only to

leading order in α , which is sufficient for our purpose here and furthermore simplifies the calculations considerably. The inclusion of QED $\mathcal{O}(\alpha)$ corrections modifies the QCD evolution equation for charged parton distributions by an additional term which, in an obvious symbolic notation, reads [15, 16]

$$\dot{q}(x, Q^2) = \frac{\alpha_s}{2\pi} [P_{qq} * q + P_{qg} * g] + \frac{\alpha}{2\pi} e_q^2 P_{qq}^\gamma * q \quad (4)$$

with $P_{qq}^\gamma(z) = \left(\frac{1+z^2}{1-z}\right)_+$. Notice that the addition [12, 14] of a further term $(\alpha/2\pi) e_q^2 P_{q\gamma} * \gamma$ to the r.h.s. of eq. (4) would actually amount to a subleading $\mathcal{O}(\alpha^2)$ contribution since the photon distribution $\gamma(x, Q^2)$ of the nucleon is of $\mathcal{O}(\alpha)$ [17, 18, 19, 20, 21, 22]. The standard QCD evolution equation for the gluon distribution remains unaltered at leading $\mathcal{O}(\alpha)$. The perturbative (p) $\mathcal{O}(\alpha)$ modifications of the valence quark distributions in (3), as implied by (4), are now given by

$$\begin{aligned} \delta_p u_v(x, Q^2) &= \frac{\alpha}{2\pi} \int_{Q_0^2}^{Q^2} d \ln q^2 \int_x^1 \frac{dy}{y} P\left(\frac{x}{y}\right) u_v(y, q^2) \\ \delta_p d_v(x, Q^2) &= -\frac{\alpha}{2\pi} \int_{Q_0^2}^{Q^2} d \ln q^2 \int_x^1 \frac{dy}{y} P\left(\frac{x}{y}\right) d_v(y, q^2) \end{aligned} \quad (5)$$

with $P(z) = (e_u^2 - e_d^2)P_{qq}^\gamma(z)$ and $Q_0^2 \equiv \mu_{\text{LO}}^2 = 0.26 \text{ GeV}^2$, $u_v(x, q^2)$, $d_v(x, q^2)$ are taken from the dynamical (radiative) parton model [13]. Notice that this treatment of the perturbative components differs from that in [12].

For the nonperturbative (np) modifications we estimate, following eq. (12) in [12],

$$\begin{aligned} \delta_{np} u_v(x) &= \frac{\alpha}{2\pi} \ln \frac{Q_0^2}{\mu_0^2} \int_x^1 \frac{dy}{y} P\left(\frac{x}{y}\right) u_v(y, Q_0^2) \\ \delta_{np} d_v(x) &= -\frac{\alpha}{2\pi} \ln \frac{Q_0^2}{\mu_0^2} \int_x^1 \frac{dy}{y} P\left(\frac{x}{y}\right) d_v(y, Q_0^2) \end{aligned} \quad (6)$$

where we take $\mu_0 = m_q \simeq 10 \text{ MeV}$, i.e. of the order of the current quark masses [2]. Here the parton distributions at $Q^2 \leq Q_0^2$ were taken to equal their values at the perturbative input scale $Q^2 = Q_0^2$, i.e. were ‘frozen’.

In fig. 1 we show the total $\delta_p q_v(x, Q^2) + \delta_{np} q_v(x)$ and the purely perturbative $\delta_p q_v(x, Q^2)$ isospin violating distributions at a typical scale of $Q^2 = 10 \text{ GeV}^2$. Our total

isospin violating distributions are quite similar to those in [12], as well as to those in [5] which were obtained by rather different methods. The fact that only our total results are compatible with nonperturbative (bag) model expectations [5, 9, 10, 11] is indicative for the necessity of our nonperturbative estimates in (6) and that the rather marginal perturbative contributions in (5) are not sufficient for a realistic estimate of isospin violating effects. It is, furthermore, particularly interesting to note that our results depend, as mentioned above, only on a single free parameter, i.e. μ_0 in (6).

Encouraged by this agreement we present in fig. 2 and 3 the corresponding predictions for the isospin violating distribution $\delta\bar{u}$ and $\delta\bar{d}$ of sea quarks as defined in (3) and obtained from (5) and (6) by replacing u_v and d_v by $\bar{u}$ and $\bar{d}$, respectively. At $Q^2 = 10 \text{ GeV}^2$ the perturbative contribution $\delta_p\bar{q}(x, Q^2)$ in fig. 2 does not dominate the total result $\delta_p\bar{q}(x, Q^2) + \delta_{np}\bar{q}(x, Q^2)$ which is obviously in contrast to the predictions at higher scales as illustrated, for example, at $Q^2 = M_W^2$ in fig. 3. Similar results are obtained for the LO CTEQ4 parton distributions [23] where also valence-like sea distributions are employed at an input scale $Q_0^2 = 0.49 \text{ GeV}^2$, i.e., $x\bar{q}(x, Q_0^2) \rightarrow 0$ as $x \rightarrow 0$. Such predictions may be tested by dedicated precision measurements of Drell–Yan and DIS processes employing neutron (deuteron) targets as well.

As a possible immediate application of our predictions for the isospin violating valence distributions let us finally turn to the relation in (1). Since the isospin violation generated by the QED $\mathcal{O}(\alpha)$ correction is such as to remove more momentum from up–quarks than down–quarks, as is evident from fig. 1, it works in the right direction to reduce the NuTeV anomaly [1], i.e. $\sin^2\theta_W = 0.2277 \pm 0.0013 \pm 0.0009$ as compared to the world average of other measurements [2] $\sin^2\theta_W = 0.2228(4)$. This effect is also slightly Q^2 dependent because of the perturbative contribution in (5). At $Q^2 = 10 \text{ GeV}^2$, appropriate for the NuTeV experiment, our total distributions in fig. 1 imply $\delta U_v = -0.002226$ and $\delta D_v = 0.000890$ which, together with $U_v + D_v = 0.3648$ in (2), leads to a change in the measured value of $\sin^2\theta_W$ of $\delta\sin^2\theta_W = \delta R_I^- = -0.0020$ according to (1). Since the value

of $\sin^2 \theta_W$ from the NuTeV experiment, with $\delta R_I^- \equiv 0$ in (1), is 0.0049 larger than the world average of other measurements, our predicted isospin violation δR_I^- reduces this discrepancy by about 40%. This reduction may be slightly diminished if it is corrected for experimental acceptance cuts [3].

To summarize, we evaluated the modification of parton distributions due to QED leading $\mathcal{O}(\alpha)$ corrections to the standard QCD evolution equations. For the isospin violating valence $\delta q_v(x, Q^2)$ and sea $\delta \bar{q}(x, Q^2)$ distributions ($q = u, d$) unique perturbative predictions are obtained within the dynamical (radiative) parton model. The nonperturbative contributions to δq_v and $\delta \bar{q}$ have been estimated and depend on a single free parameter chosen to be the current quark mass. The results for the valence distributions bear some similarity to nonperturbative bag-model expectations. Our total predictions for δq_v reduce significantly the discrepancy between the large result for $\sin^2 \theta_W$ as derived from deep inelastic $\nu(\bar{\nu})N$ data (‘NuTeV anomaly’) and the world average of other measurements.

References

- [1] G.P. Zeller et al., NuTeV Collab., *Phys. Rev. Lett.* **88** (2002) 091802
- [2] Particle Data Group, S. Eidelman et al., *Phys. Lett.* **B592** (2004) 1
- [3] G.P. Zeller et al., NuTeV Collab., *Phys. Rev.* **D65** (2002) 111103(R), **D67** (2003) 119902(E), and references therein
- [4] J.T. Londergan and A.W. Thomas, *Phys. Rev.* **D67** (2003) 111901(R)
- [5] J.T. Londergan and A.W. Thomas, *Phys. Lett.* **B558** (2003) 132, and references therein
- [6] S. Davidson et al., *JHEP* **02** (2002) 037
- [7] S. Kretzer et al., *Phys. Rev. Lett.* **93** (2004) 041802

- [8] E.A. Paschos and L. Wolfenstein, *Phys. Rev.* **D7** (1973) 91
- [9] E. Sather, *Phys. Lett.* **B274** (1992) 433
- [10] E.N. Rodionov, A.W. Thomas, J.T. Londergan, *Mod. Phys. Lett.* **A9** (1994) 1799
- [11] C.J. Benesh and J.T. Londergan, *Phys. Rev.* **C58** (1998) 1218
- [12] A. D. Martin et al., *Eur. Phys. J.* **C39** (2005) 155
- [13] M. Glück, E. Reya, A. Vogt, *Eur. Phys. J.* **C5** (1998) 461, and references therein
- [14] M. Roth and S. Weinzierl, *Phys. Lett.* **B590** (2004) 190
- [15] J. Kripfganz and H. Perlt, *Z. Phys.* **C41** (1988) 319
- [16] H. Spiesberger, *Phys. Rev.* **D52** (1995) 4936
- [17] J. Blümlein, G. Levman, H. Spiesberger, *J. Phys.* **G19** (1993) 1695
- [18] M. Glück, M. Stratmann, W. Vogelsang, *Phys. Lett.* **B343** (1995) 399
- [19] B. Kniehl, *Phys. Lett.* **B254** (1991) 267
- [20] A. De Rujula, W. Vogelsang, *Phys. Lett.*, **B451** (1999) 437
- [21] M. Glück, C. Pisano, E. Reya, *Phys. Lett.* **B540** (2002) 75
- [22] V. Lenderman, H.C. Schultz–Coulon, D. Wegener, *Eur. Phys. J.* **C31** (2003) 343,
and references therein
- [23] H.L. Lai et al., CTEQ4, *Phys. Rev.* **D55** (1997) 1280




Figure 1: The isospin violating ‘majority’ δu_v and ‘minority’ δd_v valence quark distributions at $Q^2 = 10 \text{ GeV}^2$ as defined in (3). The perturbative GRV(pert.) predictions are calculated according to (5). With the non-perturbative contribution in (6) added, one obtains the total estimates GRV(total). The bag model estimates are taken from ref. [5].




Figure 2: The isospin violating sea distributions $\delta\bar{u}$ and $\delta\bar{d}$ at $Q^2 = 10 \text{ GeV}^2$ as defined in (3) with u_v, d_v replaced by $\bar{u}, \bar{d}$. The perturbative (pert.) predictions are calculated according to (5) with u_v, d_v replaced by $\bar{u}, \bar{d}$. The same replacement holds for (6) when calculating the non-perturbative contributions which have to be added to the ‘pert.’ predictions in order to obtain the total results. The LO GRV sea distributions [13] are used throughout.




Figure 3: As in fig. 2 but at $Q^2 = M_W^2$.